

ON INSTABILITY OF RUBAKOV-SHAPOSHNIKOV MODEL

G. LAVRELASHVILI

ABSTRACT. Stability of six dimensional Rubakov-Shaposhnikov model is reinvestigated. It is shown that the model is stable with respect to tensor and vector perturbations and it is unstable in scalar perturbations sector with very particular instability pattern: there are no unstable modes for the first two lowest angular harmonics, $m = 0$ and $m = 1$, whereas there is a single negative mode for each higher m .

რეზიუმე. გამოკვლეულია რუბაკოვ-შაპოშნიკოვის ექვს განზომილებიანი მოდელის სტაბილობა. ნაჩვენებია რომ მოდელი სტაბილურია ტენზორული და ვექტორული შეშფოთების მიმართ და არასტაბილურია სკალარული შეშფოთებების სექტორში ძალიან განსაკუთრებული სქემით: პირველი ორი უდაბლესი კუთხური ჰარმონიკისათვის, $m = 0$ და $m = 1$, უარყოფითი მოდები არ არსებობენ, მაშინ როდესაც ყოველი მაღალი m -ისათვის ნაპოვნია თითო უარყოფითი მოდა.

I. INTRODUCTION

In the beginning of 80th two very important papers by Rubakov and Shaposhnikov were published in the same issue of Physics Letter B [1], [2]. The first one [1] was discussing possibility that (in modern language) we live on a brane in higher dimensional space and in the second one [2] the warped compactification was introduced in order to attack cosmological constant problem. These ideas created basis for “extra dimensional revolution” which happen 15 years later [3], [4], [5], [6], [7], [8].

The stability of the Rubakov-Shaposhnikov model with warped compactification [2] was questioned [9] soon after the model was suggested. It was found that the model is stable under tensor and vector perturbations, but has unstable modes in scalar perturbations sector. Recently we became aware that there is an algebraic error in the prove namely in the equation (23) of [9]. The aim of present note is to correct this error and reinvestigate the stability of the Rubakov-Shaposhnikov model. Since in addition there are numerous misprints in the most of equations in the journal version of

2010 *Mathematics Subject Classification.* 83E15.

Key words and phrases. Extra dimensions, cosmological constant problem, warped compactification.

[9], first we repeat here derivation of Schrödinger equations for linear perturbations and then give direct numerical proof of existence of an unstable modes in scalar perturbations sector.

II. SPONTANEOUS COMPACTIFICATION WITH ZERO COSMOLOGICAL CONSTANT

To describe the solution, leading to zero four dimensional cosmological constant, proposed in [2] we consider gravity in $d + N$ -dimensional space-time with the metric \hat{g}_{AB} (signature $+-\dots-$). The Einstein equations is written with the cosmological constant:

$$\hat{R}_{AB} - \frac{1}{2}\hat{g}_{AB}\hat{R} = \Lambda\hat{g}_{AB}. \quad (1)$$

It is assumed that $\Lambda > 0$. With the warped ansatz for the metric

$$\hat{g}_{AB} = \begin{pmatrix} \sigma(x^a)g_{\mu\nu}(x^\lambda) & 0 \\ 0 & \tilde{g}_{ab}(x^a) \end{pmatrix}, \quad (2)$$

the equation (1) reduces to the system of equations

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \Lambda_{\text{phys}}g_{\mu\nu}, \quad (3)$$

$$\tilde{R}_{ab} = -\frac{2}{N+d-2}\Lambda\tilde{g}_{ab} + d\left(\frac{\tilde{\nabla}_a\tilde{\nabla}_b\sigma}{2\sigma} - \frac{\tilde{\nabla}_a\sigma\tilde{\nabla}_b\sigma}{4\sigma^2}\right), \quad (4)$$

$$\frac{1}{2}\tilde{\nabla}_a\tilde{\nabla}^a\sigma + (d-2)\frac{\tilde{\nabla}_a\sigma\tilde{\nabla}^a\sigma}{4\sigma} - \frac{2\Lambda}{N+d-2}\sigma = -\frac{2\Lambda_{\text{phys}}}{d-2}, \quad (5)$$

where the hats and tildas respectively denote $d + N$ -dimensional and N -dimensional quantities, $\mu, \nu, \dots = 0, 1, \dots, d-1$, and $a, b, \dots = d, \dots, d+N-1$.

In these equations Λ_{phys} is an arbitrary parameter arising from the separation of variables, $R_{\mu\nu}$ is constructed from $g_{\mu\nu}$ and \tilde{R}_{ab} and $\tilde{\nabla}_a$ are constructed from \tilde{g}_{ab} according to the usual rules. The Latin indices a, b, \dots are raised and lowered with the metric \tilde{g}_{ab} . Note that, in contrast to the standard approach to spontaneous compactification, the space defined by the metric \hat{g}_{AB} is not the direct product of the d - and N -dimensional spaces. This difference is related to the presence of a warped factor $\sigma(x^a)$ in front of $g_{\mu\nu}(x)$ and is critical future in the entire discussion.

The equation (3) is the Einstein equation for the d -dimensional metric $g_{\mu\nu}$ with the cosmological constant Λ_{phys} . For $\Lambda_{\text{phys}} = 0$ it has a solution corresponding to a flat space. Assuming $d = 4$ and $N = 2$, the equations (4) and (5) can be solved with the result:

$$\tilde{g}_{ab} = \begin{pmatrix} -1 & 0 \\ 0 & f(\rho) \end{pmatrix}, \quad (6)$$

$$f(\rho) = -\frac{8}{5\Lambda} \left[\operatorname{tg}\left(\sqrt{\frac{5\Lambda}{8}}\rho\right) \right]^2 \left[\cos\left(\sqrt{\frac{5\Lambda}{8}}\rho\right) \right]^{4/5}, \quad (7)$$

$$\sigma(\rho) = \left[\cos\left(\sqrt{\frac{5\Lambda}{8}}\rho\right) \right]^{4/5}, \quad (8)$$

where $\Lambda_{\text{phys}} = 0$, $x^4 \equiv \rho$, $\rho \in [0, \rho_{\text{max}}]$, $x^5 \equiv \theta$, $\theta \in [0, 2\pi]$.

In spite of the fact that this solution is noncompact in the usual sense (the circumference $x^\mu = \text{const}$, $\rho = \text{const}$ can be arbitrarily large, $l = 2\pi\sqrt{-f} \rightarrow \infty$ as $\rho \rightarrow \rho_{\text{max}}$), it can be shown that the presence of the two extra dimensions is unobservable at low energies [2].

III. STABILITY ANALYSIS

A. The equations of motion. Let us find the equations of motion for a small fluctuations about the solution \hat{g}_{AB}^0 . For this we substitute $\hat{g}_{AB} \rightarrow \hat{g}_{AB}^0 + \epsilon_{AB}$ into the Einstein equations (1). Neglecting all terms with powers higher than first in ϵ_{AB} , we find

$$\hat{\nabla}_C \hat{\nabla}_A \epsilon_B^C + \hat{\nabla}_C \hat{\nabla}_B \epsilon_A^C - \hat{\nabla}_C \hat{\nabla}^C \epsilon_{AB} - \hat{\nabla}_A \hat{\nabla}_B \epsilon_C^C = \frac{4\Lambda}{2-N-d} \epsilon_{AB}, \quad (9)$$

where the covariant derivatives $\hat{\nabla}_A$ are calculated using the background metric \hat{g}_{AB}^0 . In what follows we set $d = 4$ and $N = 2$. Next we use the fact that \hat{g}_{AB}^0 is independent of x_μ and go to the momentum representation in x_μ . Denoting

$$\begin{aligned} \epsilon_{\mu\nu} &= \sigma h_{\mu\nu}, & \epsilon_\mu^\nu &= h_\mu^\nu, & \epsilon^{\mu\nu} &= \sigma^{-1} h^{\mu\nu}, \\ \epsilon_{a\mu} &= A_{a\mu}, & \epsilon_a^\mu &= \sigma^{-1} A_a^\mu, & \epsilon_{ab} &= \varphi_{ab}, \end{aligned} \quad (10)$$

and substituting the decomposition of $h_{\mu\nu}$ and $A_{a\mu}$ into components with spin 0, 1 and 2

$$\begin{aligned} h_{\mu\nu}(k, x^a) &= l_{\mu\nu} + k_\mu f_\nu + k_\nu f_\mu + \frac{k_\mu k_\nu}{k^2} P + \left(\eta_{\mu\nu} - \frac{k_\mu k_\nu}{k^2} \right) S, \\ A_{a\mu}(k, x^a) &= r_{a\mu} + k_\mu \Phi_a, \end{aligned} \quad (11)$$

where

$$k^\mu l_{\mu\nu} = 0, \quad k^\mu r_{a\mu} = 0, \quad k^\mu f_\mu = 0, \quad l_\mu^\mu = 0, \quad (12)$$

from the equation (9) we obtain seven separate equations for tensor, vector and scalar (under rotations of x_μ) perturbations:

$$k^2 \sigma^{-1} l_{\mu\nu} - (\tilde{\nabla}_a \tilde{\nabla}^a + 2\pi_a \tilde{\nabla}^a) l_{\mu\nu} = 0; \quad (13)$$

$$i(\tilde{\nabla}^a + \pi^a) r_{a\nu} - \sigma(\tilde{\nabla}_a \tilde{\nabla}^a + 2\pi_a \tilde{\nabla}^a) f_\nu = 0, \quad (14a)$$

$$\begin{aligned} k^2 \sigma^{-1} r_{a\mu} + ik^2 \tilde{\nabla}_a f_\mu - (\tilde{\nabla}_b \tilde{\nabla}^b + \pi_b \tilde{\nabla}^b - \Lambda) r_{a\mu} + \\ + (\tilde{\nabla}_b \tilde{\nabla}^b - \pi_a \tilde{\nabla}^b + 2\pi_b \tilde{\nabla}^a - \tilde{\nabla}_b \pi_a - 2\pi_a \pi_b) r_\mu^b = 0; \end{aligned} \quad (14b)$$

$$k^2\sigma^{-1}S + ik^2\sigma^{-1}\pi_a\Phi^a - \frac{1}{2}\pi_a\tilde{\nabla}^a(P + 3S + \varphi_b{}^b) + \\ + (2\pi^a\pi^b + \tilde{\nabla}^a\pi^b + \pi^b\tilde{\nabla}^a)\varphi_{ab} - (\tilde{\nabla}_a\tilde{\nabla}^a + 2\pi_a\tilde{\nabla}^a)S = 0, \quad (15a)$$

$$2ik^2\sigma^{-1}(\tilde{\nabla}^a + \pi^a)\Phi_a + k^2\sigma^{-1}(2S + \varphi_b{}^b) - \\ - (\tilde{\nabla}_a\tilde{\nabla}^a + 2\pi_a\tilde{\nabla}^a)(P - S) = 0, \quad (15b)$$

$$i(\tilde{\nabla}_b\varphi_a{}^b - \tilde{\nabla}_a(3S + \varphi_b{}^b) + \frac{1}{2}\pi_a\varphi_b{}^b + \pi_b\varphi_a{}^b) - (\tilde{\nabla}_b\tilde{\nabla}^b + 2\pi_b\tilde{\nabla}^b - \Lambda)\Phi_a + \\ + (\tilde{\nabla}_b\tilde{\nabla}_a - \pi_a\tilde{\nabla}_b + 2\pi_b\tilde{\nabla}_a - \tilde{\nabla}_b\pi_a - 2\pi_a\pi_b)\Phi^b = 0, \quad (15c)$$

$$k^2\sigma^{-1}\varphi_{ab} + ik^2\sigma^{-1}(\tilde{\nabla}_a\Phi_b + \tilde{\nabla}_b\Phi_a) - \frac{1}{2}(\pi_b\tilde{\nabla}_a\pi_a\tilde{\nabla}_b)(P + 3S) - \\ - \tilde{\nabla}_a\tilde{\nabla}_b(P + 3S) - \tilde{\nabla}_a\tilde{\nabla}_b\varphi_c{}^c + (\tilde{\nabla}_c + 2\pi_c)(\tilde{\nabla}_a\varphi_b{}^c + \tilde{\nabla}_b\varphi_a{}^c) + \\ + (\Lambda - \tilde{\nabla}_c\tilde{\nabla}^c - 2\pi_c\tilde{\nabla}^c)\varphi_{ab} = 0, \quad (15d)$$

where $\pi_a = (\partial/\partial x^a)\ln\sigma$.

The system of equations (13-15) is a system of eigenvalue equations with the role of the unknown eigenvalue played by k^2 . If a system (for example, (15)) is consistent, there will be at least one equation of the form

$$L\Psi(k, x^a) = k^2\Psi(k, x^a), \quad (16)$$

for some $\Psi(k, x^a)$, where L is a differential operator acting on x^a . The question of stability of the solution \hat{g}_{AB}^0 now reduces to the question of the possible values of k^2 . The solution will be linearly stable if k^2 has no negative values and solution is unstable if k^2 can take negative values.

Note that in this problem there is gauge invariance related to the invariance of the equation (1) under general coordinate transformations $x^A \rightarrow x'^A = x^A - \Delta x^A$. In terms of variables equations (10) and (11), gauge transformations with the parameters $\Delta x^A = [\sigma^{-1}(\omega^\mu + k^\mu\eta), \chi^a]$ looks like

$$\delta l_{\mu\nu} = 0, \quad \delta r_{a\mu} = \left(\frac{\partial}{\partial x^a} - \pi_a\right)\omega_\mu, \quad \delta f_\mu = i\sigma^{-1}\omega_\mu, \\ \delta\Phi_a = \left(\frac{\partial}{\partial x^a} - \pi_a\right)\eta + i\chi_a, \quad \delta S = \pi_a\chi^a, \quad (17) \\ \delta P = 2ik^2\sigma^{-1}\eta + \pi_a\chi^a, \quad \delta\varphi_{ab} = \left(\frac{\partial\tilde{g}_{ab}}{\partial x^c} + \tilde{g}_{ac}\frac{\partial}{\partial x^b} + \tilde{g}_{bc}\frac{\partial}{\partial x^a}\right)\chi^c.$$

B. The mass spectrum. It follows directly from the equation (13) for tensor perturbations that k^2 is non-negative for this sector. In order to see this, we set $\mu, \nu = 0$ in this equation, multiply both sides by $\sigma^2\sqrt{-f}l^{00}(-k) = [-\det(\hat{g}_{AB}^0)]^{1/2}l^{00}(-k)$, and integrate over $dx^4 dx^5 \equiv d\tilde{x}$. Integrating the right hand side by parts (the correctness of this procedure can be rigorously justified), we obtain the equation

$$k^2 \int \sigma\sqrt{-f}d\tilde{x}|l_{00}(k)|^2 = - \int \sigma^2\sqrt{-f}d\tilde{x}[\tilde{\nabla}_a l_{00}(k)]^* [\tilde{\nabla}^a l^{00}(k)], \quad (18)$$

from which it follows that $k^2 \geq 0$.

For vector perturbations first we have to fix the gauge. We choose the gauge conditions as

$$f_\mu = 0. \quad (19)$$

Furthermore, we set $\mu = 0$ in the second of equations (14), multiply by $\sigma\sqrt{-f}r^{a0}(-k)$, sum over a , and integrate over $d\tilde{x}$. Then with the help of gauge condition equation (19) and use of the first of equation (14) after some transformations we find

$$\begin{aligned} k^2 \int \sqrt{-f}d\tilde{x}r^{a0}r_{a0} &= \\ &= \int \sigma\sqrt{-f}d\tilde{x} \left(-\tilde{\nabla}_a r_{b0}\tilde{\nabla}^a r^{b0} - \frac{1}{2}\pi_a\pi_b r^{a0}r_0^b - \frac{3}{4}\pi_c\pi^c r^{a0}r_{a0} \right). \end{aligned} \quad (20)$$

The integral multiplying k^2 and the right hand side are both non-positive. Therefore, for the vector perturbations also $k^2 \geq 0$.

Let us now turn to equations equation (15) for scalar perturbations. We choose the gauge condition in the form

$$\Phi_a = 0, \quad P - S = 0. \quad (21)$$

We shall assume that $k^2 \neq 0$. Then equation (15a) is a consequence of the three other equations and can be omitted. From equation(15b) we obtain

$$S = -\frac{1}{2}\varphi_b{}^b, \quad (22)$$

which can be taken as a definition of S in terms of $\varphi_a{}^b$. Using this equation and the gauge condition equation (21), the two remaining equations can be written in terms of $\varphi_a{}^b$:

$$(\tilde{\nabla}_a + \pi_a)\varphi_b{}^b = -2(\tilde{\nabla}_b + \pi_b)\varphi_a{}^b, \quad (23a)$$

$$\begin{aligned} k^2\sigma^{-1}\varphi_{ab} &= -(\tilde{\nabla}_a\tilde{\nabla}_b + \pi_a\tilde{\nabla}_b + \pi_b\tilde{\nabla}_a)\varphi_c{}^c - \\ &-(\tilde{\nabla}_c + 2\pi_c)(\tilde{\nabla}_a\varphi_b{}^c + \tilde{\nabla}_b\varphi_a{}^c) + (\tilde{\nabla}_c\tilde{\nabla}^c + 2\pi_c\tilde{\nabla}^c - \Lambda)\varphi_{ab}. \end{aligned} \quad (23b)$$

Equation (23a) gives two relations between three variables φ_{ab} , so a single independent variable remains. Three equations (23b) are equivalent to each other and determine the spectrum of k^2 . The problem is to solve the constraint (23a), that is, to express all the in terms of a single independent variable ξ and its derivatives. Then from (23b) we find following equation for ξ :

$$k^2\xi = M\xi, \quad (24)$$

where M is a differential operator acting on x^a .

To proceed further let us expand $\varphi_{ab}(k, \rho, \theta)$ in a Fourier series in θ

$$\varphi_{ab}(\rho, \theta) = \sum_{m=-\infty}^{\infty} e^{im\theta}\varphi_{abm}(\rho), \quad (25)$$

and consider cases $m = 0$ and $m \neq 0$ separately.

For $m = 0$ the equation (23a) gives:

$$(\partial_\rho + \Gamma + \pi_4)\varphi_5^4 = 0, \quad (26a)$$

$$(\partial_\rho + \pi_4)(3\varphi_4^4 + \varphi_5^5) = -2\Gamma(\varphi_4^4 - \varphi_5^5), \quad (26b)$$

where $\Gamma = \tilde{\Gamma}_{45}^5 = \frac{1}{2}\partial_\rho \ln f$. The first of these equations has no regular solutions except for zero, so it gives $\varphi_5^4 = 0$. Using this condition, the equation (26b), the background equation (5) and denoting $3\varphi_4^4 + \varphi_5^5 \equiv \xi_0$, we find

$$k^2 \sigma^{-1} \xi_0 = -\partial_\rho^2 \xi_0 + \left(2\frac{\partial_\rho \Gamma}{\Gamma} - \Gamma - \pi_4\right) \partial_\rho \xi_0 + \left(2\pi_4 \frac{\partial_\rho \Gamma}{\Gamma} - 2\pi_4 \Gamma - \pi_4^2\right) \xi_0. \quad (27)$$

Multiplying this equation by ξ_0 and integrating over ρ from zero to ρ_{\max} with the weight $W(\rho)$ given by the expression

$$W(\rho) = \exp\left(-\int d\rho \left(2\frac{\partial_\rho \Gamma}{\Gamma} - \Gamma - \pi_4\right)\right) \geq 0, \quad (28)$$

after integration by parts we obtain

$$k^2 \int \sigma^{-1} W \xi_0^2 d\rho = \int W d\rho \left[(\partial_\rho \xi_0)^2 + \left(2\pi_4 \frac{\partial_\rho \Gamma}{\Gamma} - 2\pi_4 \Gamma - \pi_4^2\right) \xi_0^2 \right]. \quad (29)$$

The positivity of the last term in the integrand follows from the explicit form of background solution, (7) and (8). Therefore, $k^2 \geq 0$ for $m = 0$.

In the $m \neq 0$ case we denote $\tilde{\varphi}_5^4 m(\rho) = \frac{1}{im} \varphi_5^4 m(\rho)$. Dropping the index m on the fields φ_{ab} , we rewrite equation (23a) as

$$(\partial_\rho + \pi_4)(3\varphi_4^4 + \varphi_5^5) = 2\mu \tilde{\varphi}_5^4 - 2\Gamma(\varphi_4^4 - \varphi_5^5), \quad (30a)$$

$$\varphi_4^4 + 3\varphi_5^5 = -2(\partial_\rho + \Gamma + \pi_4)\tilde{\varphi}_5^4, \quad (30b)$$

where $\mu = -m^2/f$. The solution of this system has the form:

$$\begin{aligned} \xi &\equiv 3\varphi_4^4 + \varphi_5^5 + 2\Gamma\tilde{\varphi}_5^4, & G &\equiv \mu + \partial_\rho \Gamma, \\ \tilde{\varphi}_5^4 &= \frac{(\partial_\rho + \Gamma + \pi_4)\xi}{2G}, & 3\varphi_4^4 + \varphi_5^5 &= \xi - \Gamma \frac{(\partial_\rho + \Gamma + \pi_4)\xi}{G}, \\ \varphi_4^4 - \varphi_5^5 &= \frac{1}{2\Gamma} (2\mu\tilde{\varphi}_5^4 - (\partial_\rho + \pi_4)(3\varphi_4^4 + \varphi_5^5)). \end{aligned} \quad (31)$$

After some awkward algebra, the equation (23b) gives following for ξ :

$$k^2 \sigma^{-1} \xi = -\partial_\rho^2 \xi + A(\rho) \partial_\rho \xi + B(\rho) \xi, \quad (32)$$

$$A(\rho) = 2\frac{\partial_\rho G}{G} - \Gamma, \quad B(\rho) = \mu + \frac{1}{2}\pi_4^2 + 2\pi_4 \Gamma + 2(\Gamma + \pi_4) \frac{\partial_\rho G}{G}. \quad (33)$$

It is convenient to change to the dimensionless variable $x = \sqrt{\frac{5\Lambda}{8}}\rho$, $x \in [0, \pi/2]$, in the equations (32, 33).

Now we bring Schrödinger equation (32) into standard form in two steps. First we change independent variable from x to τ according to $d\tau = dx/\sqrt{\sigma}$. When $x \in [0, \pi/2]$ variable τ is changing from 0 to τ_{max} ,

$$\tau_{max} = \int_0^{\pi/2} \frac{dx}{[\cos(x)]^{2/5}} = \frac{\pi^{3/2} \csc(\frac{3\pi}{10})}{2\Gamma(\frac{7}{10})\Gamma(\frac{4}{5})} \approx 2.27221542. \quad (34)$$

This way we get rid of factor σ in the l.h.s. of equation (32) and coefficients A and B are changed to

$$A_\tau = \sqrt{\sigma}A + \frac{1}{2}\sqrt{\sigma}\pi_4, \quad B_\tau = \sigma B. \quad (35)$$

Next with the transformation $\xi = \exp(\frac{1}{2} \int A_\tau d\tau)\chi$ we get rid of the first derivative term and arrive at

$$\left(-\frac{d^2}{d\tau^2} + U\right)\chi = k^2\chi, \quad \text{with } U = B_\tau + \frac{1}{4}A_\tau^2 - \frac{1}{2}\partial_\tau A_\tau. \quad (36)$$

Potential U close to $\tau \rightarrow 0$ behaves as

$$U = \frac{35}{4\tau^2} + O(\tau^2) \quad \text{for } m = 1, \quad (37)$$

and

$$U = \frac{m^2 - \frac{1}{4}}{\tau^2} + \frac{56 - 8m^2}{15} + O(\tau^2) \quad \text{for } m \geq 2, \quad (38)$$

and close to τ_{max} as

$$U = -\frac{1}{4(\tau_{max} - \tau)^2}. \quad (39)$$

Note that quantum mechanical potential $U = -\gamma/x^2$ for $\gamma > \gamma_{cr} = 1/4$ corresponds to unstable situation (“falling” to the center, see e.g. [10]). So, in equation (36, 39) exactly critical case is realized, which is on a border between stability and instability. Another observation is that the $m = 1$ case is distinguished, because potential U is not negative in the inner region, while starting from $m = 2$ it is negative not only asymptotically $\tau \rightarrow \tau_{max}$, equation (39), but also in the inner region, Figure 1.

The regular branch of the wave function χ close to $\tau \rightarrow 0$ behaves as

$$\chi \propto \tau^{7/2} - \frac{k^2}{16}\tau^{11/2} \quad \text{for } m = 1, \quad (40)$$

and

$$\chi \propto \tau^{m+1/2} + \frac{56 - 8m^2 - 15k^2}{60(m+1)}\tau^{m+5/2} \quad \text{for } m \geq 2, \quad (41)$$

and close to $\tau \rightarrow \tau_{max}$ as

$$\chi \propto (\tau_{max} - \tau)^{1/2}. \quad (42)$$

To determine the number of bound states of Schrödinger equation in a given potential we investigated the zero energy wave function. According

to known theorems (see e.g. [11]) the number of nodes of zero energy wave function exactly counts the number of negative energy states in a given potential. Solving numerically Schrödinger equation (36) with above boundary conditions we found that there are no negative modes in $m = 1$ case, whereas there is single negative mode for each higher m . We checked this statement up to $m = 10$.




FIGURE 1. Shape of the potential $U(\tau)$ for $m = 1, 2$ and 5 .

IV. CONCLUDING REMARKS

We have shown that the solution described by equations (2,6,7,8) and corresponding to the $\Lambda_{\text{phys}} = 0$ is linearly unstable. The instability is related to θ -dependant perturbations, which are scalars under rotation of the four dimensional coordinates x_μ . We found that a single unstable mode appears in spectrum of linear perturbations for each angular harmonic with $m \geq 2$. A similar situation can be expected to arise for small Λ_{phys} . Even if the solution is stable starting from some $\Lambda_{\text{phys}}^0 \neq 0$, it is quite improbable that the value of Λ_{phys}^0 will be $\sim 10^{-56} \text{cm}^{-2}$, in agreement with current observations [12], [13].

Although the Rubakov-Shaposhnikov solution is found to be classically unstable, knowledge of unstable modes can be useful, since it suggests the form of the stable solution to be sought. Since the perturbations leading to instability are asymmetric under θ -rotations, it is clear that the initially symmetric state of the system tends to the more favored asymmetric state. Lorentz invariance is not violated in the development of the instability, as

it would be if the instability were related to the vector perturbations. So, it is quite possible that the equations (4) and (5) for $N = 2$ have asymmetric, θ -dependent solution, which might be stable.

Since Rubakov-Shaposhnikov model is basic ingredient for many modern higher dimensional setups it is natural to ask whether the instability disappears by adding extra fields. So, question of stability should be carefully checked in each case.

ACKNOWLEDGEMENTS

I would like to thank Valery Rubakov for valuable comments on the manuscript. The main part of work has been done during my visits to Geneva University, Switzerland, the Albert-Einstein-Institute, Golm, Germany and the Abdus Salam ICTP, Trieste, Italy. I would like to thank the theory groups of these institutions and especially Ruth Durrer, Hermann Nicolai and Seif Randjbar-Daemi for kind hospitality. I also thank the Tomalla foundation and Georgian National Science Foundation (Grant #GNSF/ST08/4 – 405) for the financial support and deeply acknowledge the ICTP associate membership program. I am grateful to Zurab Ratiani for pointing out an algebraic error in the Eq. (23) of Ref. [9].

REFERENCES

1. V. A. Rubakov and M. E. Shaposhnikov, Do We Live Inside A Domain Wall? *Phys. Lett. B* **125** (1983), 136–138.
2. V. A. Rubakov and M. E. Shaposhnikov, Extra Space-Time Dimensions: Towards A Solution To The Cosmological Constant Problem. *Phys. Lett. B* **125** (1983), 139–143.
3. N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, The hierarchy problem and new dimensions at a millimeter. *Phys. Lett. B* **429** (1998), 263–272. [arXiv:hep-ph/9803315].
4. I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, New dimensions at a millimeter to a Fermi and superstrings at a TeV. *Phys. Lett. B* **436** (1998), 257–263. [arXiv:hep-ph/9804398].
5. N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phenomenology, astrophysics and cosmology of theories with sub-millimeter dimensions and TeV scale quantum gravity. *Phys. Rev. D* **59** (1999), 086004 [arXiv:hep-ph/9807344].
6. M. Gogberashvili, Hierarchy problem in the shell-universe model. *Int. J. Mod. Phys. D* **11** (2002), 1635–1638. [arXiv:hep-ph/9812296].
7. L. Randall and R. Sundrum, A large mass hierarchy from a small extra dimension. *Phys. Rev. Lett.* **83** (1999), 3370–3373. [arXiv:hep-ph/9905221].
8. L. Randall and R. Sundrum, An alternative to compactification. *Phys. Rev. Lett.* **83** (1999), 4690–4693. [arXiv:hep-th/9906064].
9. G. Lavrelashvili and P. G. Tinyakov, The possibility of spontaneous compactification leading to zero cosmological constant. (Russian) *translated from Yadernaya Fiz.* **41** (1985), No. 1, 271–277, *Soviet J. Nuclear Phys.* **41** (1985), No. 1, 171–177, *Yad. Fiz.* **41** (1985) 272–275.
10. L. D. Landau and E. M. Lifshitz, Quantum Mechanics, V3 of Course of Theoretical Physics. *Butterworth-Heinemann*, 1998.

11. H. Amann and P. Quittner, A nodal theorem for coupled systems of Schrodinger equations and the number of bound states. *J. Math. Phys.* **36** (1995), No. 9, 4553–4560.
12. S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], Measurements of Omega and Lambda from 42 High-Redshift Supernovae. *Astrophys. J.* **517** (1999), 565–586. [arXiv:astro-ph/9812133].
13. A. G. Riess *et al.* [Supernova Search Team Collaboration], Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *Astron. J.* **116** (1998), 1009–1038. [arXiv:astro-ph/9805201].

(Received 13.01.2012)

Author's address:

A. Razmadze Mathematical Institute
I. Javakhishvili Tbilisi State University
2, University Str., Tbilisi 0186
Georgia
E-mail: lavrela@itp.unibe.ch