




Ministry of Communication and Information
Technologies of Azerbaijan Republic



the Abdus Salam International
Centre for Theoretical Physics



Institute of Cybernetics
of Azerbaijan National
Academy of Sciences



Baku State University



Institute of Information
Technology of Azerbaijan
National Academy of Sciences

PCI'2008

**THE SECOND INTERNATIONAL CONFERENCE
"PROBLEMS OF CYBERNETICS AND INFORMATICS"**

Dedicated to the 50th Anniversary of the ICT in Azerbaijan

Volume II


September 10-12, 2008
Baku, Azerbaijan



Ministry of Communication and Information
Technologies of Azerbaijan Republic



the Abdus Salam International
Centre for Theoretical Physics



Institute of Cybernetics
of Azerbaijan National
Academy of Sciences



Baku State University



Institute of Information
Technology of Azerbaijan
National Academy of Sciences

PCI'2008

THE SECOND INTERNATIONAL CONFERENCE
"PROBLEMS OF CYBERNETICS AND INFORMATICS"
Dedicated to the 50th Anniversary of the ICT in Azerbaijan

Volume II

September 10-12, 2008
Baku, Azerbaijan

<i>Gambar Guluyev, Asif Rzayev, Farhad Pashayev, R.Malikov, Y.Aliyev, M.Rizvanov.</i>	176
“Technologist” local control system of technological processes.....	
<i>Mamed Akhmedov, J. Mamedov, Ch. Gheydarov.</i> Application of intelligence models for functional analyze of technological units	180
<i>Rouzbeh Rahmanian.</i> DEA: a method for measuring the efficiency of agencies	184
<i>Alovsat Aliyev, Roza Shahverdiyeva.</i> On need for working out the models and mechanisms of establishment and management of ICT-Technoparks	188
<i>Irina Gorodnitsky.</i> Evaluation of derivative time-delay modeling for robust pitch detection in very high and nonstationary noise	192
<i>Vasily Makarov, Tahir Alizada.</i> Algorithm of estimation of evolving models parameters ..	196
<i>Irina Gorodnitsky, Tahir Alizada, Sevinj Guliyeva, Dilara Mastaliyeva, Ulkar Rzayeva.</i> Fourier-based method for estimating signal perturbations in linearly-correlated noise	200
<i>Tahir Alizada, Ulviyya Mammadova, Elchin Aliyev, Rauf Gadimov.</i> Identification of false impulses in the binary components of noisy signal	204
Section #4. Applied Stochastic Analysis.....	207
<i>M. Antónia Amaral Turkman, Elisete Correia.</i> Bayes inference on the ratio between two proportions.....	209
<i>Che Soong Kim, Agassi Melikov, Vagif Feyziyev.</i> Analysis of multirate systems with shared reservation of channels and queues of wideband calls.....	210
<i>Vladimir Korolyuk.</i> Problems of singular perturbations in analysis of stochastic systems ...	214
<i>Zurab Kvatadze, Tengiz Shervashidze.</i> Some limit theorems for I.I.D. and conditionally independent random variables	217
<i>Anvar Norjigitov, Olimjon Sharipov.</i> On the central limit theorem for weakly dependent random variables with values in $D[0,1]$	220
<i>Jakhongir Azimov.</i> Asymptotic behaviors of the critical branching processes with decreasing immigration	223
<i>Tamilla Nasirova, Elshan Ibayev.</i> Fourier transformation with respect to phase of the conditional distribution of one of the semi-Markovian random walk processes	225
<i>Ruslan Meilanov, Vladimir Zaborovsky, Vetlugin Beibolaev, Ramazan Magomedov.</i> Fractal analysis of network processes.....	229
<i>Ruslan Meilanov, Vladimir Zaborovsky, Vladimir Mulukha.</i> Models and characteristics of TCP flows in stochastic network environment.....	232
<i>Yakubdján Khusanbaev.</i> On functional limit theorem for stochastic branching processes with immigration	236
<i>Elizbar Nadaraya, Petre Babilua.</i> On the power of the goodness-of-fit test based on wolverton-wagner density estimates.....	238
<i>Omar Purtukhia, Vakhtang Jaoshvili.</i> Stochastic integral representation of Poisson functionals	242
<i>Petre Babilua, Besarion Dochviri.</i> On the pricing of a European option.....	246
<i>Zurab Zerakidze.</i> Estimation of parameters of chi-square statistical structures	249
<i>Omar Glonti, Zaza Khechinashvili.</i> Martingale measures for the geometrical Gaussian martingale	251
<i>Grigol Sokhadze.</i> On infinit dimensional elasticity theory	253

STOCHASTIC INTEGRAL REPRESENTATION OF POISSON FUNCTIONALS

Omar Purtukhia, Vakhtang Jaoshvili

Ivane Javakhishvili Tbilisi State University,
 A. Razmadze Mathematical Institute Georgian National Academy of Science
 Georgia, e-mail: o.purtukhia@math.sci.tsu.ge

0. In the theory of random processes special place take the so-called martingale representation theorems which, for example, implies the representation of the Wiener and Poisson functionals in the form of stochastic integrals. In the 80th of the past century, it turned out (see Harison and Pliska (1981)) that the martingale representation theorems (along with the Girsanov's measure change theorem) play an important role in the modern financial mathematics. In particular, using the integrand of the stochastic integral appearing in the integral representation, one can construct hedging strategies in the European options of different type.

According to the well-known result obtained by Clark (1970), if F is a \mathfrak{F}_T -measurable random variable with $EF^2 < \infty$, then there exists the adapted process $\varphi_t(\omega) \in L_2([0, T] \times \Omega)$, such that the integral representation

$$F = EF + \int_0^T \varphi_t(\omega) dw_t \quad (P\text{-a.s.})$$

holds. However, this result says nothing on finding the process $\varphi_t(\omega)$ explicitly. In this direction we are familiar with one sufficiently general result, the so-called Ocone-Clark's formula by which for the Wiener functionals: $\varphi_t(\omega) = E[D_t^w F | \mathfrak{F}_t^w](\omega)$, where $D_t^w F$ is the stochastic derivative (so-called the Malliavin's derivative) of the functional F . Another distinct method of finding an integrand $\varphi_t(\omega)$ belongs to Shyryaev, Yor (2003), when the functional ξ is of "maximal" type. With the functional they linked the associated Lewy's martingale and used the generalized Ito's formula. According to the this result, if $S_t = \max_{u \leq t} w_u$, then the following integral representation is holds:

$$S_T = ES_T + 2 \int_0^T [1 - \Phi(\frac{S_t - w_t}{\sqrt{T-t}})] dw_t,$$

where $\Phi(\cdot)$ is the standard normal distribution function.

One should note that application of the Ocone-Clark formula needs as a rule, on the one hand, essential efforts, and, on the other hand, in the cases if the functional F has no the stochastic derivative, its application is impossible. Our approach within the classical Ito's calculus allows one to construct $\varphi_t(\omega)$ explicitly by using both the standard L_2 theory and the theories of weighted Sobolev spaces, if the functional F has no stochastic derivative (see Jaoshvili, Purtukhia (2005)). It is known that the events indicator has, in general, no stochastic derivative: more exactly, the indicator function of $A \in \mathfrak{F}_T^w$ belongs to $D_{2,1}^w$ (where $D_{2,1}^w$ denotes the space of square integrable Wiener functionals having the first order stochastic derivative) if and only if $P(A)$ equal to zero or one. Consequently, one cannot apply the Ocone-Clark formula for the indicator $I_{\{w_T > 0\}}$ ($P(w_T > 0) = 1/2$), whereas our approach allows one to write the following representation:

$$I_{\{w_T > 0\}} = EI_{\{w_T > 0\}} + \int_0^T \Phi_{0, T-t}(w_t) dw_t,$$

where $\Phi_{0, T-t}(\cdot)$ is the function of a normal distribution with parameters 0 and $T-t$. The subsequent generalization of the Ocone-Clark formula to the so-called normal martingales (the martingale is said to be normal, if $\langle M, M \rangle_t = t$) is due to Ma, Protter, Martin (1998). According to this formula, if $F \in D_{2,1}^M$, then the Ocone-Haussmann-Clark's representation

$$F = E(F) + \int_0^T {}^p(D_t^M F) dM_t$$

is valid; here $D_{2,1}^M$ denotes the space of quadratically integrable functionals having the first order stochastic derivative, and ${}^p(D_t^M F)$ is the predictable projection of the stochastic derivative $D_t^M F$ of the functional F . But, in this case (exactly, when the quadratic variation $[M, M]$ is not deterministic), unlike the Wiener's one (see, example from Ma, Protter, Martin (1998)), it is impossible to define in a generally adopted manner an operator of stochastic differentiation to obtain the structure of Sobolev spaces, which allows one to construct explicitly the stochastic derivative operator in many cases. In reality, the example from Ma, Protter, Martin (1998) shows that two definitions, Sobolev space and chaos expansion, are compatible if and only if $[M, M]$ is deterministic. Therefore in the martingale case the space $D_{p,1}^M$ ($1 < p < 2$) cannot be defined in the usual way -- i.e., by closing the class of smooth functionals with respect to the corresponding norm. In work of Purtukhia (2003) the space $D_{p,1}^M$ ($1 < p < 2$) is proposed for a class of normal martingales and the integral representation formula of Ocone-Haussmann-Clark is established for functionals from this space.

Since, the Compensated Poisson process belongs to a class of normal martingales -- $\langle M, M \rangle_t = t$, but its quadratic variation is not deterministic -- $[M, M]_t = M_t + t$, consequently, the Ocone-Haussmann-Clark's formula makes it impossible to construct explicitly the operator of the stochastic derivative of the functionals of the Compensated Poisson process, saying nothing on the construction of its predictable projection. Our approach within the framework of nonanticipative stochastic calculus of semimartingales allows one to construct explicitly the expression for the integrand of the stochastic integral in the theorem of martingale representation for square integrable functionals of the Compensated Poisson process, and to derive the formula allowing one to construct explicitly predictable projections of their stochastic derivatives (see Jaoshvili, Purtukhia (2007)).

1. Let $(\Omega, \mathfrak{F}, P, (\mathfrak{F}_t)_{0 \leq t \leq \infty})$ be a filtered probability space satisfying the usual conditions. Assume that the standard Poisson process N_t is given on it $P(N_t = n) = e^{-t} t^n / n!$, $n = 0, 1, 2, \dots$ and that \mathfrak{F}_t is generated by N ($\mathfrak{F}_t = \mathfrak{F}_t^N$), $\mathfrak{F} = \mathfrak{F}_T$. Denote $M_t := N_t - t$. Let $Z^+ = \{0, 1, 2, \dots\}$ and $P = \{P_1, P_2, P_3, \dots\}$ -- be the Poisson distribution: $P_x = e^{-T} T^x / x!$, $x = 0, 1, 2, \dots$. Let us denote $\Delta f(x) = f(x) - f(x-1)$ ($f(x) := 0$, if $x < 0$) and define the Poisson-Charle's polynomials: $\Pi_n(x) = [(-1)^n \Delta^n P_x] / P_x$, $n \geq 1$; $\Pi_0 = 1$.

It is wellknown from the course of Functional Analysis that the sequence $\{\pi_n(x)\}_{n \geq 0}$ ($\pi_n(x) = \Pi_n(x) / c_n$) is a basis in the space $L_2(Z^+)$ ($L_2(Z^+) = \{f : \sum_{x=0}^{\infty} f^2(x) < \infty\}$).

Let $\rho(x, T) := e^{-T} T^x / x!$ and denote by $L_2^T := L_2(Z^+; \rho(x, T))$ the functional space on Z^+ with the finite norm $\|g\|_{2, T} := \|g\rho^{1/2}(T)\|_{L_2}$.

Proposition 1.1. The space L_2^T is a Banach space with basis $\{x^n \rho(x, T), n \geq 1\}$.

Proposition 1.2. If $F(\cdot - T) \in L_2^T$, then the stochastic integral $\int_0^T E[F(M_T) | \mathfrak{F}_{t-}] dM_t$,

is well defined.

Let us denote $\nabla_x f(x) := f(x+1) - f(x)$, $(\nabla_x F(M_T) := \nabla_x F(x)|_{x=M_T})$.

Theorem 1.1. If $F(\cdot) \in L_2^T$ and for some number $0 < \alpha < 1$: $\nabla_x F(\cdot - T) \in L_2^{T/\alpha}$, then the stochastic integral below is well defined and the following representation is valid:

$$F(M_T) = E[F(M_T)] + \int_0^T E[\nabla_x F(M_T) | \mathfrak{F}_{t-}] dM_t \quad (P\text{-a.s.}) \quad (1.1)$$

Theorem 1.2. In the conditions of the Theorem 1.1 the following relation holds:

$${}^p[D_t^M F(M_T)] = E[\nabla_x F(M_T) | \mathfrak{F}_{t-}] \quad (dP \otimes ds\text{-a.s.}), \quad (1.2)$$

where ${}^p[D_t^M F(M_T)]$ denotes the predictable projection of the stochastic derivative (with respect to the Compensated Poisson process) $D_t^M F(M_T)$ of the functional $F(M_T)$.

Example 2.1. The random variable M_T^2 has the following stochastic integral representation:

$$M_T^2 = E[M_T^2] + \int_0^T (1 + 2M_{t-}) dM_t \quad (P\text{-a.s.})$$

(note that in the Wiener process cases the Ocone-Clark's formula gives us that:

$$w_T^2 = E[w_T^2] + \int_0^T 2w_t dw_t \quad (P\text{-a.s.}).$$

2. Fix now $0 \leq S \leq T \leq \infty$ and let us denote

$$\rho(x, y, S, T) = \frac{S^x}{x!} \cdot e^{-S} \cdot \frac{T^{y-x}}{(y-x)!} \cdot e^{-(T-S)}.$$

Denote by $L_2^{S, T} := L_2(Z^+ \times Z^+; \rho(x, y, S, T))$ the functional space on $Z^+ \times Z^+$ with the finite norm $\|g\|_{2, S, T} := \|g\rho^{1/2}(S, T)\|_{L_2}$.

Proposition 2.1. The space $L_2^{S, T}$ is a Banach space with basis $\{x^n y^m \rho(x, y, S, T), n, m \geq 1\}$.

For any function of two variables $g(\cdot, \cdot)$ introduce the designation:

$$\nabla^2 g(x, y) = g(x+1, y+1) - g(x, y).$$

It is not difficult to see that

$$\nabla_x [\nabla_y g(x, y)] = \nabla_y [\nabla_x g(x, y)]$$

and

$$\nabla^2 g(x, y) = \nabla_x [\nabla_y g(x, y)] + \nabla_x g(x, y) + \nabla_y g(x, y).$$

Let us denote

$$\begin{aligned} \nabla_i^2 g(M_S, M_T) &= \nabla_x [\nabla_y g(M_S, M_T)] I_{[0, S]}(t) I_{[0, T]}(t) + \\ &+ \nabla_x g(M_S, M_T) I_{[0, S]}(t) + \nabla_y g(M_S, M_T) I_{[0, T]}(t). \end{aligned}$$

It is clear that for any function of two variables $g(\cdot, \cdot)$ there exists the function $h(\cdot, \cdot)$, such that $g(x, y) = h(x, y - x)$. Let us denote $\bar{g}(x, y) = g(x - S, y - x - (T - S))$.

Theorem 2.1. If $F(\cdot, \cdot) \in L_2^{S, T}$ and for some number $0 < \alpha < 1$: $\nabla_x [\nabla_y \bar{F}(\cdot, \cdot)] \in L_2^{S/\alpha, T/\alpha}$, then the stochastic integral below is well defined and the following stochastic integral representation is valid:

$$F(M_S, M_T) = E[F(M_S, M_T)] + \int_0^T E[\nabla_i^2 F(M_S, M_T) | \mathfrak{F}_{i-}] dM_i \quad (P\text{-a.s.}) \quad (2.1)$$

Theorem 2.2. In the conditions of the Theorem 2.1 the following relation holds:

$${}^p[D_i^M F(M_S, M_T)] = E[\nabla_i^2 F(M_S, M_T) | \mathfrak{F}_{i-}] \quad (dP \otimes ds\text{-a.s.}), \quad (2.2)$$

where ${}^p[D_i^M F(M_S, M_T)]$ denotes the predictable projection of the stochastic derivative (with respect to the Compensated Poisson process) $D_i^M F(M_S, M_T)$ of the functional $F(M_S, M_T)$.

*The work has been financed by the Georgian National Science Foundation grant
 № 337/07, 06_223_3-104.*

References

1. Clark J. M. The representation of functionals of Brownian motion as stochastic integrals, *Annals of Mathematical Statistics* 41 (1970), 1282-1295.
2. Haussmann U. G. On the integral representation of functionals of Ito processes, *Stochastics* 3 (1979), 17-28.
3. Ocone D. Malliavin's calculus and stochastic integral representation of functionals of diffusion processes, *Stochastics* 12 (1984), 161-185.
4. Harrison J. M., Pliska S. R. Martingales and stochastic integrals in the theory of continuous trading. *Stochastic Processes and Applications* 11 (1981), 215-260.
5. Ma J., Protter P., Martin J.S. Anticipating integrals for a class of martingales. *Bernoulli* 4 (1998), 81-114.
6. Shyryaev A., Yor M. About stochastic integral representation of functionals of Brownian Motions I. *Probab. Th. And Appl.* vol. 48 (2003), 375-385.
7. Purtukhia O. An Extension of the Ocone-Haussmann-Clark Formula for a Class of Normal Martingales. *Proceedings of A. Razmadze Mathematical Institute*, vol. 132 (2003), 127-136.
8. Jaoshvili V., Purtukhia O. Stochastic Integral Representation of Functionals of Wiener Processes. *Bull. Georg. Acad. Sci.* Vol. 171, No 1 (2005), 17-20.
9. Jaoshvili V., Purtukhia O. Stochastic Integral Representation of Functionals of Poisson Processes. *Proceedings of A. Razmadze Mathematical Institute*, vol. 143 (2007), 37-60.